

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.**



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁷ : A61K	A2	(11) International Publication Number: WO 00/41669
		(43) International Publication Date: 20 July 2000 (20.07.00)
<p>(21) International Application Number: PCT/GB00/00099</p> <p>(22) International Filing Date: 14 January 2000 (14.01.00)</p> <p>(30) Priority Data: 9900752.8 15 January 1999 (15.01.99) GB</p> <p>(71) Applicant (for all designated States except US): ANGIOGENE PHARMACEUTICALS LTD. [GB/GB]; 14 Plowden Park, Aston Rowant, Watlington, Oxfordshire OX9 5SW (GB).</p> <p>(72) Inventor; and (75) Inventor/Applicant (for US only): DAVIS, Peter, David [GB/GB]; 10 Aston Park, Aston Rowant, Watlington OX9 5SW (GB).</p> <p>(74) Agents: BAILLIE, Iain, C. et al.; Langner Parry, 52-54 High Holborn, London WC1V 6RR (GB).</p>		<p>(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).</p> <p>Published Without international search report and to be republished upon receipt of that report.</p>
(54) Title: BENZIMIDAZOLE VASCULAR DAMAGING AGENTS		
<div style="text-align: center;"> <p>(I)</p> </div>		
<p>(57) Abstract</p> <p>A group of vascular damaging agents which can be used in the preparation of medicaments for the treatment of diseases involving neovascularisation are provided. These are 5(6)-substituted benzimidazole-2-carbamates of formula (I) wherein Alk is an alkyl group, X is oxygen, sulphur, sulphinyl, sulphonyl, carbonyl (CO), thiocarbonyl (CS), sulphonyloxy, NH, iminomethylene (C=NH), N-hydroxyiminomethylene, N-alkoxyiminomethylene, dialkoxymethylene, 1,3-dioxolan-2-yl, 1,1-ethenyl, a group CHR³ or a bond, R¹ is hydrogen, alkylaminocarbonyl or alkoxycarbonyl, R² is hydrogen, alkoxycarbonyl, cyanomethyl, cyanoethyl, alkoxymethyl or acetoxymethyl. R³ is hydrogen, hydroxy, alkoxy or amino, A is an optionally substituted aromatic, optionally substituted heteroaromatic, optionally substituted heterocycloalkyl, optionally substituted alkyl or optionally substituted cycloalkyl group and the pharmaceutically acceptable salts, solvates and hydrates thereof. Most of the compounds of this group are novel, in particular those in which A is an aromatic or heteroaromatic ring with substituents, particularly substituents which are phosphates or alkylphosphates. The invention therefore provides both novel compounds and pharmacological compositions with compounds within the broad definition.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

BENZIMIDAZOLE VASCULAR DAMAGING AGENTS

This invention relates to vascular damaging agents and particularly to the use of new and known substituted benzimidazoles in the preparation of medicaments for the
5 treatment of diseases involving neovascularisation.


Formation of new vasculature by angiogenesis is a key pathological feature of several diseases (J Folkman, New England Journal of Medicine 333, 1757-1763 (1995)). For example, for a solid tumour to grow it must develop its own blood supply upon which
10 it depends critically for the provision of oxygen and nutrients; if this blood supply is mechanically shut off the tumour undergoes necrotic death. Neovascularisation is also a clinical feature of skin lesions in psoriasis, of the invasive pannus in the joints of rheumatoid arthritis patients and of atherosclerotic plaques. Retinal neovascularisation is pathological in macular degeneration and in diabetic retinopathy. In all these diseases
15 reversal of neovascularisation by damaging the newly-formed vascular endothelium is expected to have a beneficial therapeutic effect.

5(6)-Substituted benzimidazole-carbamates are known and have found use as antiparasitic agents (P. J. Islip in Burgers Medicinal Chemistry (M. E. Wolff ed.),
20 Fourth Edition, Part II, p481, (1979)). Examples of such compounds include mebendazole, fenbendazole, oxibendazole, flubendazole, albendazole, cyclobendazole, parbendazole, dribendazole, luxabendazole, and etibendazole. Their mode of action for their antiparasitic action is believed to involve selective binding to tubulin of the target parasite while having little effect due to binding tubulin of the mammalian host
25 (Biochim. Biophys. Acta 630, 271-278, (1980)). Some of these compounds have been shown to be antimitotic for cancer cells and one particular 5(6)-substituted benzimidazole-2-carbamate, nocodazole, has therefore been studied as an anticancer agent (Cancer Research, 36, 905-916 (1976)). No effects on neovasculature have been reported for any of these compounds.

Some structurally-unrelated compounds which bind tubulin have been shown to have anti-vascular effects when given at their maximum tolerated dose (MTD) (S. A. Hill et al. Eur. J Cancer, 29A, 1320-1324 (1993)) but other tubulin-binding agents, such as docetaxel, have no vascular-damaging activity even when administered at the MTD.

- 5 The presence of tubulin-binding properties is then not predictive for antivascular activity.

According to the present invention there is provided the use of 5(6)-substituted benzimidazole-2-carbamates for the preparation of compositions for the treatment of
10 diseases involving angiogenesis in which the 5(6)-substituted benzimidazole carbamate has the formula



I

15

wherein

Alk is an alkyl group

X is oxygen, sulphur, sulphinyl, sulphonyl, carbonyl (CO), thiocarbonyl (CS),
20 sulphonyloxy, NH, iminomethylene (C=NH), N-hydroxyiminomethylene, N-alkoxyiminomethylene, dialkoxymethylene, 1,3-dioxolan-2-yl, 1,1-ethenyl, a group CHR³ or a bond

R¹ is hydrogen, alkylaminocarbonyl or alkoxycarbonyl

R² is hydrogen, alkoxycarbonyl, cyanomethyl, cyanoethyl, alkoxymethyl or
25 acetoxymethyl.


R³ is hydrogen, hydroxy, alkoxy or amino

A is an optionally substituted aromatic, optionally substituted heteroaromatic, optionally substituted heterocycloalkyl, optionally substituted alkyl or optionally substituted cycloalkyl group

30

	X	A
Fenbendazole	S	Ph
Mebendazole	CO	Ph
Albendazole	S	nPr
Oxibendazole	O	nPr
Nocodazole	CO	2-thienyl

Certain of these compounds are novel. In one embodiment the novel compounds are those of formula I in which at least one of the substituents on the group A is a group Y
 5 where Y is as hereinbefore defined. Particularly preferred are compounds defined by the formula



10 wherein

alk is an alkyl group

B is an aromatic or heteroaromatic ring

X is oxygen, sulphur, sulphinyl, sulphonyl, carbonyl (CO), thiocarbonyl (CS),
 15 sulphonyloxy, NH, iminomethylene (C=NH), N-hydroxyiminomethylene, N-alkoxyiminomethylene, dialkoxymethylene, 1,3-dioxolan-2yl, 1,1-ethenyl, a group CHR³ or a bond

R¹ is hydrogen, alkylaminocarbonyl or alkoxycarbonyl

R² is hydrogen, alkoxycarbonyl, cyanomethyl, cyanoethyl, alkoxymethyl or
 20 acetoxymethyl

R³ is hydrogen, hydroxy, alkoxy or amino

Y is as hereinbefore defined

R⁷ and R⁸ are each independently H, alkyl, halogen, hydroxy, amino, alkylamino, dialkylamino, alkoxy, alkylthio, cyano, nitro, or trifluoromethyl

25

with the proviso that Y is not NHC(O)Me and when B is a thiophene ring then Y is not C(O)CF_3 and when B is a 5(6)-benzimidazole ring then Y is not NHCO_2Me or NHCO_2Et

- 5 and the pharmaceutically acceptable salts, solvates, hydrates and prodrugs thereof.

As used herein the term "alkyl", alone or in combinations, means a straight or branched-chain alkyl group containing from one to seven, preferably a maximum of four, carbon atoms such as methyl, ethyl, propyl, isopropyl, butyl, sec-butyl, t-butyl
10 and pentyl. Examples of alkoxy groups are methoxy, ethoxy, propoxy, isopropoxy, butoxy and t-butoxy. The term "halogen" means fluorine, chlorine, bromine or iodine.

The term "aryl" as used herein unless otherwise stated includes reference to a C_{6-10} aryl group which may, if desired, carry one or more substituents selected from halogeno,
15 alkyl, haloalkyl, alkoxy, hydroxy, amino, nitro and cyano. The term "aralkoxy" means an alkoxy group substituted with an aryl group.

The term heteroaryl is defined herein as a mono- or bi-cyclic aromatic group containing one to four heteroatoms selected in any combination from N, S or O atoms
20 and a maximum of 9 carbon atoms. Examples of heteroaryl groups include pyridyl, pyrimidyl, furyl, thienyl, pyrrolyl, pyrazolyl, indolyl, benzofuryl, benzothienyl, benzothiazolyl, oxazolyl, isoxazolyl, thiazolyl, isothiazolyl, imidazolyl, triazolyl, quinolyl and isoquinolyl groups.

25 The term heterocycloalkyl includes heterocycloalkyl groups containing 3-6 carbon atoms and one or two oxygen, sulphur or nitrogen atoms. Particular examples of such groups include azetidiny, pyrrolidiny, piperidiny, homopiperidiny, piperaziny, homopiperaziny, morpholiny or thiomorpholiny groups.

30 The term cycloalkyl means a cycloaliphatic group containing 3-10 carbon atoms such as, for example, cyclopropyl, cyclobutyl, cyclopentyl or cyclohexyl.

One particularly preferred group of compounds are those of formula II in which Y is a phosphate group.

- 5 Another particularly preferred group of compounds are those of formula II in which Y is a group $\text{NR}^5\text{C}(\text{O})\text{R}^4$, R^5 is hydrogen and R^4 is a 1-aminoalkyl group which can be further substituted for example by a hydroxy group.


Where one or more functional groups in compounds of formula I or II are sufficiently
10 basic or acidic the formation of salts is possible. Suitable salts include pharmaceutically acceptable salts for example acid addition salts including hydrochlorides, hydrobromides, phosphates, sulphates, hydrogen sulphates, alkylsulphonates, arylsulphonates, acetates, benzoates, citrates, maleates, fumarates, succinates, lactates and tartrates, salts derived from inorganic bases including alkali metal salts such as
15 sodium or potassium salts, alkaline earth metal salts such as magnesium or calcium salts, and salts derived from organic amines such as morpholine, piperidine or dimethylamine salts.

Those skilled in the art will recognise that compounds of formulae I and II may exist as
20 stereoisomers and/or geometrical isomers and accordingly the present invention includes all such isomers and mixtures thereof. The substituted benzimidazole group is capable of existing in tautomeric forms and the formulae I and II are intended to represent all tautomeric forms and the substituent AX- is in the 5(6) position.

25 Compounds of Formula I or II may be prepared by any process known to a person skilled in the art. Compounds of Formula I or II may be prepared by a number of processes as generally described below and more specifically in the Examples hereinafter. In the following process description, the symbols A, X and alk when used in the formulae depicted are to be understood to represent those groups described
30 above in relation to formula I or II unless otherwise indicated. In the schemes described below it may be necessary to employ protecting groups which are then

removed during the final stages of the synthesis. The appropriate use of such protecting groups and processes for their removal will be readily apparent to those skilled in the art.

- 5 Thus according to a further aspect of the invention compounds of formulae I and II in which R^1 and R^2 are hydrogen may be prepared by treatment of a diamine of formula III with a 1,3-bis(alkoxycarbonyl)-S-alkyl isothiurea, for example 1,3-bis(methoxycarbonyl)-S-methyl isothiurea or 1,3-bis(ethoxycarbonyl)-S-methyl isothiurea, in a solvent such as an alcohol, for
10 example methanol or ethanol, optionally mixed with water, at from about room temperature to the reflux temperature of the solvent for about 5 minutes to 6 hours. The reaction medium is preferably made acidic by the addition of for example an organic acid such as acetic acid or p-toluenesulphonic acid.



Compounds of formula III are either known or can be prepared by conventional procedures.

20

Compounds of formulae I and II can also be prepared from other compounds of formulae I and II by chemical modification. Example of such chemical modifications that may be applied are standard alkylation, acylation, reduction, oxidation, sulphation, aromatic halogenation, aromatic nitration, phosphorylation, hydrolysis, condensation,
25 cleavage and coupling reactions. These reactions may be used to add new substituents, to modify existing substituents or to modify other parts of the molecule.

Thus for example a compound of formula I or II in which R^1 is hydrogen can be converted into the corresponding compounds where R^1 is alkylaminocarbonyl by

treatment with an alkyl isocyanate in a solvent such as tetrahydrofuran at a temperature in the range 0° to 40°C, typically room temperature.

5 In another general example a thioether group in a compound of formula I or II can be converted into a sulphinyl group by treatment with periodate in an aqueous alcohol solvent such as aqueous methanol or in aqueous acetonitrile at about -20° to 50°C, for about 1 to 16 h. Alternatively this conversion can be effected by treatment with one equivalent of a peracid such as 3-chloroperbenzoic acid in a chlorinated solvent such as dichloromethane or chloroform, at a temperature of about -30°C to room temperature.

10

In a further general example a thioether group in a compound of formula I or II can be converted into a sulphonyl group by treatment with two or more equivalents of a peracid such as 3-chloroperbenzoic acid in a chlorinated solvent such as dichloromethane or chloroform, at a temperature of about -30°C to room temperature.

15

In a further general example a keto group in a compound of formula I or II can be converted into a carbinol group by treatment with a reducing agent for example a hydride such as lithium aluminium hydride in an ether solvent such as diethyl ether or tetrahydrofuran at a temperature of from about 0° to the reflux temperature of the solvent.

20

In a further general example a keto group in a compound of formula I or II can be converted into an imine by treatment with ammonia in an alcoholic solvent such as ethanol at around room temperature for an extended period, for example three weeks.

25

In a further general example a keto group in a compound of formula I or II can be converted into an oxime by treatment with hydroxylamine in an alcoholic solvent such as ethanol at around room temperature to around the reflux temperature of the solvent.

30 In a further general example a compound of formula I or II containing a hydroxy group can be converted into the corresponding dihydrogenphosphate ester by treatment with

for example di-tert-butyl diethylphosphoramidite in the presence of a suitable catalyst for example tetrazole in a solvent such as an ether solvent for example tetrahydrofuran at a temperature in the range -40 to 40°C, conveniently at or near room temperature, followed by treatment with an oxidising agent for example 3-chloroperoxy benzoic acid or magnesium monoperoxyphthalate at a temperature in the range -78°C to 40°C preferably -65 to -10°C. The resulting intermediate phosphate triester is treated with an acid for example trifluoroacetic acid in a solvent such as a chlorinated solvent e.g. dichloromethane at a temperature in the range -30 to 40°C conveniently at or near 0°C to give the compound of formula I or II containing a dihydrogenphosphate ester.

10

In another general example an O-alkyl group may be cleaved to the corresponding alcohol (OH) by reaction with boron tribromide in a solvent such as a chlorinated solvent e.g. dichloromethane at a low temperature e.g. around -78°C.

15 In a further general example compounds of formula I or II may be alkylated by reaction with a suitable alkylating agent such as an alkyl halide, an alkyl toluenesulphonate, an alkyl methanesulphonate or an alkyl triflate. The alkylation reaction can be carried out in the presence of a base for example an inorganic base such as a carbonate e.g. caesium or potassium carbonate, a hydride such as sodium hydride or an alkoxide such as potassium t-butoxide in a suitable solvent such as an aprotic solvent e.g. dimethylformamide or an ether solvent such as tetrahydrofuran at a temperature of around -10 to 80°C.

25 In a further general example compounds of formula I or II containing an amine group may be acylated by treatment with a carboxylic acid and a coupling reagent, for example dicyclohexylcarbodiimide, in a suitable solvent for example an aprotic solvent such as dimethylformamide, an ether solvent such as tetrahydrofuran, a chlorinated solvent for example dichloromethane or a solvent mixture at a temperature in the range 0° to 60°, preferably about room temperature.

30

In a further general example a compound of formula I or II containing an OH group can be converted into a carbamate by reaction with an alkyl isocyanate or a carbamoyl chloride in an aprotic solvent such as dimethylformamide, an ether solvent such as tetrahydrofuran, a chlorinated solvent for example dichloromethane or a solvent mixture in the presence of a base such as a tertiary amine base for example triethylamine at a temperature in the range -20° to the reflux temperature of the solvent, conveniently at or around room temperature.

In a further general example a compound of formula I or II containing an amino group can be converted into a urea by reaction with an isocyanate or a carbamoyl chloride in an aprotic solvent such as dimethylformamide, an ether solvent such as tetrahydrofuran, a chlorinated solvent for example dichloromethane or a solvent mixture in the presence of a base such as a tertiary amine base for example triethylamine at a temperature in the range -20°C to the reflux temperature of the solvent, conveniently at or around room temperature.

In a further general example a compound of formula I or II containing a hydroxy group can be converted into a carbonate by reaction with an chloroformate in an aprotic solvent such as dimethylformamide, an ether solvent such as tetrahydrofuran, a chlorinated solvent for example dichloromethane or a solvent mixture in the presence of a base such as a tertiary amine base for example triethylamine at a temperature in the range -20°C to the reflux temperature of the solvent, preferably at or around 0°C.

Preparation of a compound of formula I or II as a single enantiomer or, where appropriate, diastereomer may be effected by synthesis from an enantiomerically pure starting material or intermediate or by resolution of the final product in a conventional manner.

Acid addition salts of the compounds of formula I or II are prepared in a conventional manner by treating a solution or suspension of the free base I or II with about one

equivalent of a pharmaceutically acceptable acid. Salts of compounds of formula I or II derived from inorganic or organic bases are prepared in a conventional manner by treating a solution or suspension of the free acid I or II with about one equivalent of a pharmaceutically acceptable organic or inorganic base. Alternatively both acid addition salts and salts derived from bases may be prepared by treatment of the parent compound with the appropriate ion-exchange resin in a standard fashion. Conventional concentration and recrystallisation techniques are employed in isolating the salts.

Compounds according to the invention are able to destroy vasculature that has been newly formed, for example tumour vasculature, while leaving unaffected normal, mature vasculature. The ability of the compounds to act in this way may be determined by the tests described hereinafter.

The compounds according to the invention are thus of particular use in the prophylaxis and treatment of cancers involving solid tumours and in the prophylaxis and treatment of diseases where inappropriate angiogenesis occurs such as diabetic retinopathy, psoriasis, rheumatoid arthritis, atherosclerosis and macular degeneration.

The compounds of the invention may be administered as a sole therapy or in combination with other treatments. For the treatment of solid tumours compounds of the invention may be administered in combination with radiotherapy or in combination with other anti-tumour substances for example those selected from mitotic inhibitors, for example vinblastine, paclitaxel and docetaxel; alkylating agents, for example cisplatin, carboplatin and cyclophosphamide; antimetabolites, for example 5-fluorouracil, cytosine arabinoside and hydroxyurea; intercalating agents for example adriamycin and bleomycin; enzymes, for example asparaginase; topoisomerase inhibitors for example etoposide, topotecan and irinotecan; thymidylate synthase inhibitors for example raltitrexed; biological response modifiers for example interferon; antibodies for example edrecolomab and antibodies against the EGFr, HER2 receptor or VEGF receptor; and anti-hormones for example tamoxifen. Such combination

treatment may involve simultaneous or sequential application of the individual components of the treatment.

For the prophylaxis and treatment of disease the compounds according to the invention
5 may be administered as pharmaceutical compositions selected with regard to the
intended route of administration and standard pharmaceutical practice. Such
pharmaceutical compositions may take a form suitable for oral, buccal, nasal, topical,
rectal or parenteral administration and may be prepared in a conventional manner
using conventional excipients. For example for oral administration the pharmaceutical
10 compositions may take the form of tablets or capsules. For nasal administration or
administration by inhalation the compounds may be conveniently delivered as a powder
or in solution. Topical administration may be as an ointment or cream and rectal
administration may be as a suppository. For parenteral injection (including intravenous,
subcutaneous, intramuscular, intravascular or infusion) the composition may take the
15 form of, for example, a sterile solution, suspension or emulsion.

The dose of a compound of the invention required for the prophylaxis or treatment of a
particular condition will vary depending on the compound chosen, the route of
administration, the form and severity of the condition and whether the compound is to
20 be administered alone or in combination with another drug. Thus the precise dose will
be determined by the administering physician but in general daily dosages may be in the
range 0.001 to 100mg/kg preferably 0.1 to 50mg/kg.

BIOLOGICAL ACTIVITY

25

The following test was used to demonstrate the activity and selectivity of compounds according to the invention.

Activity against tumour vasculature measured by fluorescent dye.

The following experiment demonstrates the ability of the compounds to damage tumour vasculature.

Tumour functional vascular volume in CaNT tumour-bearing mice was measured using the fluorescent dye Hoechst 33342 according to the method of Smith *et al* (Brit J Cancer 57, 247-253, 1988). The fluorescent dye was dissolved in saline at 6.25 mg/ml and injected intravenously at 10 mg/kg 6 hours or 24 hours after intraperitoneal drug treatment. One minute later, animals were killed and tumours excised and frozen; 10 µm sections were cut at 3 different levels and observed under UV illumination using an Olympus microscope equipped with epifluorescence. Blood vessels were identified by their fluorescent outlines and vascular volume was quantified using a point scoring system based on that described by Chalkley, (J Natl Cancer Inst, 4, 47-53, 1943). All estimates were based on counting a minimum of 100 fields from sections cut at the 3 different levels. Results are expressed as percentage reduction in vascular volume compared to control.

The activity of compounds of the invention in this assay is shown in Table 1.

Table 1: Reduction in tumour vascular volume measured by fluorescent dye

Compound	Dose (mg/kg) (i.p.)	Time (h)	% Reduction in vascular volume
Fenbendazole	500	6	44
Mebendazole	500	6	56
Albendazole	500	6	51
Oxibendazole	100	6	43
Nocodazole	100	24	23
Compound of Example:			
1	500	6	81
2	500	6	80
3	50	24	22
4	50	24	53
5	50	24	99
12	50	24	51
14	50	24	39

15	50	24	61
16	50	24	56
17	50	24	62
18	50	24	47
19	50	24	88
20	50	24	69
21	50	24	74

The following non-limiting Examples illustrate the invention:

5 Example 1

Methyl [5(6)-(4-hydroxybenzoyl)-1H-benzimidazol-2yl]carbamate.

A solution of 3,4-diamino-4'-hydroxybenzophenone (49.6mg, 0.21mmol) and 1,3-bis(methoxycarbonyl)-S-methyl isothiurea (98mg, 0.44mmol) in ethanol (2.5ml) was treated with p-toluenesulphonic acid (7mg) and the mixture heated at reflux for 10 minutes. The mixture was cooled and the precipitate collected by filtration and washed with ethanol and hexane to give the title compound (16mg) as a white solid m.p. >258°C, ¹H-NMR (400 MHz, d₆-DMSO) δ 7.80 (s, 1H), 7.65 (d, 2H, J=8Hz), 7.50 (s, 1H), 7.49 (s, 1H), 6.89 (d, 2H, J=8Hz), 3.77 (s, 3H) ppm. m/e 311 (M⁺). Anal. Calculated for C₁₆H₁₃N₃O₄: C, 61.73; H, 4.21; N 13.49. Found: C, 61.68; H, 4.18; N 13.36.

Example 2

Methyl [5(6)-(4-phosphonooxybenzoyl)-1H-benzimidazol-2yl]carbamate.

A solution of methyl [5-(4-(di-tert-butylphosphonooxy)benzoyl)-1H-benzimidazol-2yl]carbamate (1.5g, 3.0mmol) in dichloromethane (55ml) was cooled in an ice bath and treated with trifluoroacetic acid (6ml) dropwise. The mixture was allowed to warm to room temperature and stirred for 1 hour before solvents were removed under reduced pressure. The residue was triturated with ether to give the title compound (1.13g) as a white solid m.p. >258°C, ¹H-NMR (300 MHz, d₆-DMSO) δ: 7.84 (s, 1H), 7.75 (d, 1H, J=9Hz), 7.57 (d, 1H, J=8Hz), 7.52 (d, J=8Hz), 7.33 (d, 1H, J=9Hz), 3.78 (s, 3H) ppm.

The methyl [5(6)-(4-(di-*tert*-butylphosphonooxy)benzoyl)-1*H*-benzimidazol-2yl]carbamate used as starting material was prepared as follows:

A solution of methyl [5(6)-(4-hydroxybenzoyl)-1*H*-benzimidazol-2yl]carbamate (100mg, 0.3mmol) in anhydrous tetrahydrofuran (1ml) stirred under a nitrogen atmosphere was treated with di-*tert*-butyl diethylphosphoramidite (74mg, 0.29mmol) and 1*H*-tetrazole (54mg, 0.78mmol) and the mixture stirred until the reaction was shown to be complete by TLC (about 1h). The cooled (-40°C) mixture was treated with 3-chloroperbenzoic acid (79mg, 0.39mmol) in dichloromethane (1ml) and stirred 10minutes before being allowed to warm to room temperature. The mixture was washed with saturated aqueous sodium bicarbonate followed by brine and the dried (MgSO₄) organic phase concentrated under reduced pressure. The residue was chromatographed on silica gel eluting first with 4% methanol/dichloromethane then with 5% methanol/dichloromethane. Methyl [5(6)-(4-(di-*tert*-butylphosphonooxy)benzoyl)-1*H*-benzimidazol-2yl]carbamate (35mg) was obtained as a white solid m.p. 99-101°C.

Example 3

Methyl [5(6)-(4-phosphonooxyphenylthio)-1*H*-benzimidazol-2yl]carbamate

A solution of methyl [5(6)-(4-(di-*tert*-butylphosphonooxy)phenylthio)-1*H*-benzimidazol-2yl]carbamate (180mg) in anhydrous dichloromethane (10ml) at 4°C was treated with trifluoroacetic acid (1ml) and stirred for 1.5h. The mixture was allowed to warm to room temperature and concentrated under reduced pressure. Ethyl acetate was added and the mixture concentrated again. The residue was triturated with diethyl ether, washed with diethyl ether followed by acetone/water 9:1 and dried to give the title compound (134mg) as a white solid m.p. 190-193°C. Anal. Calculated for C₁₅H₁₄N₃O₆PS.3H₂O: C, 40.1; H, 4.5; N 9.3. Found: C, 40.1; H, 3.9; N 9.2.

The methyl [5(6)-(4-(di-*tert*-butylphosphonooxy)phenylthio)-1*H*-benzimidazol-2yl]carbamate used as starting material was prepared as follows:

A solution of methyl [5-(4-hydroxyphenylthio)-1*H*-benzimidazol-2yl]carbamate (200mg) in a mixture of anhydrous dimethylformamide (2ml) and anhydrous

tetrahydrofuran (2ml) was treated with di-*tert*-butyl diethylphosphoramidite (350mg) and the mixture stirred for 48h at room temperature. The mixture was cooled to -65°C and treated gradually with magnesium monoperoxyphthalate (850mg) so that the temperature remained below -50°C. A saturated aqueous solution of sodium bicarbonate was added, keeping the temperature below -40°C during the addition then allowing the mixture to warm to room temperature. The mixture was extracted with three portions (50ml each) of ethyl acetate and the combined extracts washed with brine (50ml), dried (MgSO₄) and concentrated under reduced pressure. The residue was purified by radial chromatography on silica gel eluting with dichloromethane/methanol 9:1 to give methyl [5(6)-(4-(di-*tert*-butylphosphonooxy)phenylthio)-1*H*-benzimidazol-2-yl]carbamate (180mg) as a white foam.

Example 4

15 Methyl [5(6)-(4-aminophenylthio)-1*H*-benzimidazol-2-yl]carbamate

Methyl [5(6)-(4-(acetylamino)phenylthio)-1*H*-benzimidazol-2-yl]carbamate (602 mg, 1.78 mmol) was dissolved in mixture of methanol (24 ml) and hydrochloric acid (10%, 6 ml) and heated under reflux for 16h. The solution was neutralised with ammonia solution and the methanol removed under reduced pressure. The white precipitate was collected by filtration, washed with water and dried in vacuo to give 392 mg of a pale yellow solid m.p. 282-284°C. m/e 298 (M⁺).

Example 5

Methyl [5(6)-(4-alanylaminophenylthio)-1*H*-benzimidazol-2-yl]carbamate

25 A suspension of methyl [5(6)-(4-(N α -*tert*-butoxycarbonylalanyl)amino)phenylthio)-1*H*-benzimidazol-2-yl]carbamate (250mg) in dichloromethane (20ml) was treated with trifluoroacetic acid (4ml). The mixture was allowed to warm to room temperature and concentrated under reduced pressure. Ethyl acetate was added and the mixture concentrated again. The residue was triturated with diethyl ether to afford the trifluoroacetic acid salt of the title compound (105mg) as a white solid m.p. 178-

182°C. m/e 485 (M⁺). Anal. Calculated for C₁₈H₁₉N₅O₃S.2C₂HF₃O₂ C; 43.1, H; 3.5, N; 11.4 Found C; 42.8, H; 3.8, N; 11.3

The methyl [5(6)-(4-(N α -*tert*-butoxycarbonylalaninylamino)phenylthio)-1*H*-benzimidazol-2yl]carbamate used as starting material was prepared as follows:

- 5 A suspension of methyl [5(6)-(4-aminophenylthio)-1*H*-benzimidazol-2yl]carbamate (150mg) in anhydrous tetrahydrofuran (4ml) was treated with N-*tert*-butoxycarbonylalanine (100mg), cooled to -35°C and treated with 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide (100mg). The mixture was allowed to warm to room temperature and stir for 16h. Water (40ml) and ethyl acetate (20ml)
- 10 were added and the mixture extracted with four portions of ethyl acetate (50ml each). The combined extracts were washed with brine (50ml), dried (MgSO₄) and concentrated under reduced pressure. The residue was purified on silica gel eluting with ethyl acetate to give methyl [5(6)-(4-(N α -*tert*-butoxycarbonylalaninylamino)phenylthio)-1*H*-benzimidazol-2yl]carbamate
- 15 (258mg) as a white solid m.p. 222-224°C. m/e 485 (M⁺).

Example 6

Methyl [5(6)-(4-(methoxycarbonylamino)phenylthio)-1*H*-benzimidazol-2yl]carbamate

- Methyl [5(6)-(4-aminophenylthio)-1*H*-benzimidazol-2yl]carbamate (150 mg, 0.48
- 20 mmol) was dissolved in dimethylformamide (2 ml) and tetrahydrofuran (2 ml) and methyl chloroformate (40 mg, 0.42 mmol) added together with triethylamine (43 mg, 0.42 mmol). The solution was stirred at 20°C for 72 h and then evaporated to dryness under reduced pressure. The residue was purified by chromatography on silica gel eluting with 5 % methanol/dichloromethane to give the title compound as a white
- 25 solid; mp >350°C (dec.).

Example 7

Methyl [5(6)-(4-(phenylaminocarbonylamino)phenylthio)-1*H*-benzimidazol-2yl]carbamate

- 30 Methyl [5(6)-(4-aminophenylthio)-1*H*-benzimidazol-2yl]carbamate (150 mg, 0.48 mmol) was dissolved in dimethylformamide (2 ml) and tetrahydrofuran (2 ml) and

phenylisocyanate (56.8 mg, 0.48 mmol) added together with triethylamine (50 mg, ca.0.5 mmol). The solution was stirred at 20°C for 12 h and the solvents removed in vacuo. The residue was purified on silica (ethyl acetate/hexane, 2:1) to give the title compound as a white solid; mp 335–340°C (dec.).m/e 433 (M^+).

5

Example 8

Methyl [5(6)-(4-(methoxycarbonyloxy)benzoyl)-1*H*-benzimidazol-2yl]carbamate

Methyl [5(6)-(4-hydroxybenzoyl)-1*H*-benzimidazol-2yl]carbamate (110 mg, 0.35 mmol) was dissolved in dimethylformamide (3.5 ml) and triethylamine (0.5 ml). The solution was cooled to 0°C and methyl chloroformate (50 mg, 0.52 mmol) added with stirring. The solution was stirred for 0.5 h at 0°C and the 1 h at 20°C and evaporated to dryness. The residue was dissolved in ethyl acetate (50 ml) and washed with sodium bicarbonate (sat., aq., 50 ml) and brine (50 ml), dried and evaporated. The residue was purified by radial chromatography on silica gel eluting with ethyl acetate/hexane, 1:1 followed by ethyl acetate to give the title compound as a white solid; mp 224–226°C (dec.).m/e 369 (M^+).

15

Prepared in an analogous fashion to Example 1 were:

20 Example 9

Methyl [5(6)-(2-methoxycarbonylphenylthio)-1*H*-benzimidazol-2yl]carbamate from methyl 2-(3,4-diaminophenylthio)benzoate (1.5g) and 1,3-bis(methoxycarbonyl)-*S*-methyl isothiurea (2.25g) there was obtained the title compound (1.51g) as a white solid m.p. 228-230 m/e 357 (M^+). Anal. Calculated for $C_{17}H_{15}N_3O_4S$: C, 57.13; H, 4.23; N 11.75. Found: C, 57.39; H, 4.13; N 11.81.

25

Example 10

Methyl [5(6)-(3-methoxycarbonylphenylthio)-1*H*-benzimidazol-2yl]carbamate from methyl 3-(3,4-diaminophenylthio)benzoate (246mg) and 1,3-bis(methoxycarbonyl)-*S*-methyl isothiurea (371mg) there was obtained the title compound (183mg) as an off-

30

white solid m.p. 226-228 m/e 357 (M⁺). Anal. Calculated for C₁₇H₁₅N₃O₄S: C, 57.13; H, 4.23; N 11.75. Found: C, 56.58; H, 4.31; N 11.86.

Example 11

- 5 Methyl [5(6)-(4-methoxycarbonylphenylthio)-1H-benzimidazol-2-yl]carbamate from methyl 4-(3,4-diaminophenylthio)benzoate (830mg) and 1,3-bis(methoxycarbonyl)-S-methyl isothiurea (1.25g) there was obtained the title compound (694mg) as an off-white solid m.p. 280-282 m/e 357 (M⁺). Anal. Calculated for C₁₇H₁₅N₃O₄S: C, 57.13; H, 4.23; N 11.75. Found: C, 57.21; H, 4.31; N 11.73.

10

Example 12

- Methyl [5(6)-(4-hydroxyphenylthio)-1H-benzimidazol-2-yl]carbamate from 4-(4-hydroxyphenylthio)-1,2-phenylenediamine (4g) and 1,3-bis(methoxycarbonyl)-S-methyl isothiurea (6g) there was obtained the title compound (2.4g) as a white solid
15 m.p. 202-204°C. m/e 315 (M⁺). ¹H-NMR (400 MHz, d₆-DMSO) δ 3.74 (s, 3H), 6.76 and 7.2 (AA'BB', 4H, J = 8.6 Hz), 7.03 (dd, 1H, J = 1.7, 6.6 Hz), 7.29 (d, 1H, J = 1.4 Hz), 7.34 (d, 1H, J = 8.3 Hz), 9.68 (b, 1H), 11.67 (b, 2H) ppm.

Example 13

- 20 Methyl [5(6)-(4-(acetylamino)phenoxy)-1H-benzimidazol-2-yl]carbamate from 4-(4-(acetylamino)phenoxy)-1,2-phenylenediamine (1.02g) and 1,3-bis(methoxycarbonyl)-S-methyl isothiurea (2.67g) there was obtained the title compound (0.93g) as a white solid m.p. 304-306°C m/e 340 (M⁺). Anal. Calculated for C₁₇H₁₆N₄O₄: C, 60.00; H, 4.74; N 16.46. Found: C, 60.03; H, 4.72; N 16.42.

25

Prepared in an analogous fashion to Example 3 was:

Example 14

- Methyl [5(6)-(4-phosphonooxyphenoxy)-1H-benzimidazol-2-yl]carbamate from methyl
30 [5(6)-(4-(di-*tert*-butylphosphonooxy)phenoxy)-1H-benzimidazol-2-yl]carbamate

(200mg) there was obtained the title compound (140mg) as a white solid m.p. 272-275°C.

Example 15

- 5 Methyl [5(6)-(4-hydroxy- α -hydroxyiminobenzyl)-1*H*-benzimidazol-2yl]carbamate
[5(6)-(4-hydroxybenzoyl)-1*H*-benzimidazol-2yl]carbamate (100 mg, 0.32 mmol) was added to a solution of hydroxylamine (0.8 mmol) in MeOH (20 mL, prepared by treatment of hydroxylamine hydrochloride (0.64 g, 1.6 mmol) and NaOH (0.14g, 1.6 mmol) followed by filtration). The solution was heated at 70°C for 36 h, cooled and
10 water (30 ml) added. The solution was filtered and washed with ether and the solid triturated with methanol to yield the title compound (40 mg) as a white solid; mp 288–290°C.

The following known compounds were prepared by literature methods:

15

Example 16: Methyl [5(6)-(4-(acetylamino)phenoxy)-1*H*-benzimidazol-2yl]carbamate

Example 17: Methyl [5(6)-(4-aminophenoxy)-1*H*-benzimidazol-2yl]carbamate

- 20 Example 18: Methyl [5(6)-(3-aminophenoxy)-1*H*-benzimidazol-2yl]carbamate

Example 19: [5(6)-(4-hydroxybenzoyl)-1*H*-benzimidazol-2yl]carbamate


Example 20: [5(6)-(2-hydroxybenzoyl)-1*H*-benzimidazol-2yl]carbamate

25

Example 21: [5(6)-(3-hydroxybenzoyl)-1*H*-benzimidazol-2yl]carbamate

CLAIMS

1. Use of vascular damaging agents in the preparation of compositions for the treatment of diseases involving angiogenesis, characterised in that the agents are
- 5 5(6)-substituted benzimidazole-2-carbamates of formula:

**I**

wherein

10

Alk is an alkyl group

X is oxygen, sulphur, sulphinyl, sulphonyl, carbonyl (CO), thiocarbonyl (CS), sulphonyloxy, NH, iminomethylene (C=NH), N-hydroxyiminomethylene, N-alkoxyiminomethylene, dialkoxymethylene, 1,3-dioxolan-2-yl, 1,1-ethenyl, a group

15

CHR³ or a bond

R¹ is hydrogen, alkylaminocarbonyl or alkoxycarbonyl

R² is hydrogen, alkoxycarbonyl, cyanomethyl, cyanoethyl, alkoxymethyl or acetoxymethyl.

R³ is hydrogen, hydroxy, alkoxy or amino

20

A is an optionally substituted aromatic, optionally substituted heteroaromatic, optionally substituted heterocycloalkyl, optionally substituted alkyl or optionally substituted cycloalkyl group


and the pharmaceutically acceptable salts, solvates and hydrates thereof.

25

2. A use according to claim 1 wherein the substituent on A is selected from (a) alkyl substituted by one or more of hydroxy, amino, alkylamino, dialkylamino, halogen, carboxyl, SO₃H, sulphate, phosphate, alkoxycarbonyl, aralkoxycarbonyl, alkoxycarbonylamino, aminoalkylaminocarbonyl and cyano; or halogen, hydroxy,

- amino, alkoxy, alkylthio, cyano, nitro, sulphate, isothiocyanate, aryl, heteroaryl or heterocycloalkyl; or (b) a group Y selected from phosphate, alkylphosphate, $C(O)R^4$, $OC(O)R^4$, SO_2R^4 , $NHC(O)R^4$, $NR^5C(O)R^4$, SR^4 , $S(O)R^4$, OSO_2R^4 , $NHSO_2R^4$, $NR^5SO_2R^4$, SO_3H , CO_2H and CO_2R^5 where R^4 is selected from hydrogen, R^5 , OR^5 , NHR^5 , NR^5R^6 , aryl, heteroaryl or heterocycloalkyl such aryl, heteroaryl or heterocycloalkyl groups being optionally substituted with one or more substituents selected from alkyl, heterocycloalkyl, haloalkyl, hydroxy, nitro, cyano, amino, alkylamino, dialkylamino, halogen, carboxyl, SO_3H , sulphate and phosphate wherein R^5 and R^6 , which may be the same or different, are each an alkyl group optionally substituted with one or more substituents selected from hydroxy, amino, alkylamino, dialkylamino, guanidino, halogen, carboxyl, SO_3H , sulphate, phosphate, aryl and heteroaryl.

3. A 5(6)-substituted benzimidazole-2-carbamate of formula



I

wherein

20

Alk is an alkyl group

X is oxygen, sulphur, sulphinyl, sulphonyl, carbonyl (CO), thiocarbonyl (CS), sulphonyloxy, NH, iminomethylene ($C=NH$), N-hydroxyiminomethylene, N-alkoxyiminomethylene, dialkoxymethylene, 1,3-dioxolan-2-yl, 1,1-ethenyl, a group

25 CHR^3 or a bond

R^1 is hydrogen, alkylaminocarbonyl or alkoxy carbonyl


R^2 is hydrogen, alkoxy carbonyl, cyanomethyl, cyanoethyl, alkoxy methyl or acetoxymethyl.

R^3 is hydrogen, hydroxy, alkoxy or amino

A is an optionally substituted aromatic, optionally substituted heteroaromatic, optionally substituted heterocycloalkyl, optionally substituted alkyl or optionally substituted cycloalkyl group

in which at least one substituent is Y selected from phosphate, alkylphosphate, $C(O)R^4$, $OC(O)R^4$, SO_2R^4 , $NHC(O)R^4$, $NR^5C(O)R^4$, SR^4 , $S(O)R^4$, OSO_2R^4 , $NHSO_2R^4$, $NR^5SO_2R^4$, SO_3H , CO_2H and CO_2R^5 where R^4 is selected from hydrogen, R^5 , OR^5 , NHR^5 , NR^5R^6 , aryl, heteroaryl or heterocycloalkyl such aryl, heteroaryl or heterocycloalkyl groups being optionally substituted with one or more substituents selected from alkyl, heterocycloalkyl, haloalkyl, hydroxy, nitro, cyano, amino, alkylamino, dialkylamino, halogen, carboxyl, SO_3H , sulphate and phosphate wherein R^5 and R^6 , which may be the same or different, are each an alkyl group optionally substituted with one or more substituents selected from hydroxy, amino, alkylamino, dialkylamino, guanidino, halogen, carboxyl, SO_3H , sulphate, phosphate, aryl and heteroaryl and the pharmaceutically acceptable salts, solvates and hydrates thereof.

4. A carbamate according to claim 3 of formula:



wherein

alk is an alkyl group

B is an aromatic or heteroaromatic ring

X is oxygen, sulphur, sulphinyl, sulphonyl, carbonyl (CO), thiocarbonyl (CS), sulphonyloxy, NH, iminomethylene (C=NH), N-hydroxyiminomethylene, N-alkoxyiminomethylene, dialkoxymethylene, 1,3-dioxolan-2yl, 1,1-ethenyl, a group CHR^3 or a bond

R^1 is hydrogen, alkylaminocarbonyl or alkoxycarbonyl

R^2 is hydrogen, alkoxycarbonyl, cyanomethyl, cyanoethyl, alkoxymethyl or acetoxymethyl

R^3 is hydrogen, hydroxy, alkoxy or amino

Y is as hereinbefore defined

- 5 R^7 and R^8 are each independently H, alkyl, halogen, hydroxy, amino, alkylamino, dialkylamino, alkoxy, alkylthio, cyano, nitro, or trifluoromethyl

with the proviso that Y is not NHC(O)Me and when B is a thiophene ring then Y is not C(O)CF_3 and when B is a 5(6)-benzimidazole ring then Y is not NHCO_2Me or

- 10 NHCO_2Et

and the pharmaceutically acceptable salts, solvates, hydrates and prodrugs thereof.

5. A carbamate according to claim 4 in which Y is a phosphate group.

15

6. A carbamate according to claim 4 in which Y is NHC(O)R^4 wherein R^4 is a 1-aminoalkyl group.

7. A composition for treatment of diseases involving angiogenesis
20 comprising at least one 5(6)-substituted benzimidazole-2-carbamate of formula I as defined in claim 1 in an amount sufficient to damage new vasculature.

8. A composition according to claim 7, wherein the carbamate is as defined in claim 2.

25

9. A composition according to claim 7, wherein the carbamate has a formula II as defined in claim 3.

30